

COVID-19 Information[Public health information \(CDC\)](#)[Research information \(NIH\)](#)[SARS-CoV-2 data \(NCBI\)](#)[Prevention and treatment information \(HHS\)](#)[Español](#)

FULL TEXT LINKS

[Electromagn Biol Med.](#) 2017;36(3):295-305. doi: 10.1080/15368378.2017.1350584.

Epub 2017 Aug 4.

Impact of radiofrequency radiation on DNA damage and antioxidants in peripheral blood lymphocytes of humans residing in the vicinity of mobile phone base stations

[Zothansiamia](#)¹, [Mary Zosangzuali](#)¹, [Miriam Lalramdinpuii](#)¹, [Ganesh Chandra Jagetia](#)¹

Affiliations

Affiliation

¹ a Department of Zoology , Cancer and Radiation Biology Laboratory, Mizoram University , Aizawl , Mizoram , India.

PMID: 28777669 DOI: [10.1080/15368378.2017.1350584](#)**Abstract**

Radiofrequency radiations (RFRs) emitted by mobile phone base stations have raised concerns on its adverse impact on humans residing in the vicinity of mobile phone base stations. Therefore, the present study was envisaged to evaluate the effect of RFR on the DNA damage and antioxidant status in cultured human peripheral blood lymphocytes (HPBLs) of individuals residing in the vicinity of mobile phone base stations and comparing it with healthy controls. The study groups matched for various demographic data including age, gender, dietary pattern, smoking habit, alcohol consumption, duration of mobile phone use and average daily mobile phone use. The RF power density of the exposed individuals was significantly higher ($p < 0.0001$) when compared to the control group. The HPBLs were cultured and the DNA damage was assessed by cytokinesis blocked micronucleus (MN) assay in the binucleate lymphocytes. The analyses of data from the exposed group ($n = 40$), residing within a perimeter of 80 m of mobile base stations, showed significantly ($p < 0.0001$) higher frequency of micronuclei when compared to the control group, residing 300 m away from the mobile base station/s. The analysis of various antioxidants in the plasma of exposed individuals revealed a significant attrition in glutathione (GSH) concentration ($p < 0.01$), activities of catalase (CAT) ($p < 0.001$) and superoxide dismutase (SOD) ($p < 0.001$) and rise in lipid peroxidation (LOO) when compared to controls. Multiple linear regression analyses revealed a significant association among reduced GSH concentration ($p < 0.05$), CAT ($p < 0.001$)

and SOD ($p < 0.001$) activities and elevated MN frequency ($p < 0.001$) and LOO ($p < 0.001$) with increasing RF power density.

Keywords: Antioxidants; genotoxicity; humans; micronucleus; power density.

Related information

[MedGen](#)

LinkOut - more resources

Full Text Sources

[Taylor & Francis](#)

Other Literature Sources

[scite Smart Citations](#)

Medical

[MedlinePlus Health Information](#)

Miscellaneous

[NCI CPTAC Assay Portal](#)